

Gas Dynamics in the Galactic Bar Region

R. Fux

Geneva Observatory, CH-1290 Sauverny, Switzerland

Many early type barred spiral galaxies exhibit strong dustlanes at the leading edges of their bar (e.g. NGC 1300, NGC 1433), which from gas velocity field measurements and numerical hydrodynamical simulations are known to be the loci of shearing shocks with velocity gaps up to 200 km s^{-1} across. As an SBbc type galaxy, the Milky Way is expected to present similar dustlanes. Self-consistent high resolution 3D N -body and SPH simulations allow us to reliably identify their gaseous traces in the Galactic CO and HI $\ell - V$ distributions. The near-side branch of these dustlanes corresponds to the connecting arm feature and lies below ($b < 0$) the Galactic plane, while the far side branch, seen nearly end-on, traces a velocity-elongated feature near $\ell = -4^\circ$ and is located above the plane. The 3-kpc and 135-km s^{-1} arms are lateral arms, i.e. inner prolongations of disc spiral arms passing round the bar and finally intersecting the dustlanes at their respective latitudes (Figs. 1a-b).

Considering that the latter intersections occur close to the major axis of the bar and within the bar ends, their spatial location can provide new constraints on the inclination angle φ and corotation radius R_{CR} of the Galactic bar. Geometrically, if ℓ_1 and ℓ_2 are the longitudes where the 3-kpc and the 135-km s^{-1} arms respectively intersect the dustlanes, and s_1 and s_2 the galactocentric distances of these intersections (Fig. 1c), then:

$$\frac{s_i}{R_o} = \frac{\sin \ell_i}{\sin(\varphi + \ell_i)} \quad (i = 1, 2), \quad \varphi = -\frac{1 + q}{\cot \ell_2 + q \cot \ell_1},$$

where $q \equiv s_1/s_2$ is an unknown asymmetry parameter. The 135-km s^{-1} arm indeed involves higher forbidden velocities than the 3-kpc arm which, according to our models, means that the former arm is closer to the Galactic centre than the latter, i.e. $q > 1$. The CO observations suggest $\ell_1 = 14.5^\circ$ and $\ell_2 = -4.5^\circ$, yielding the constraints shown in Fig. 1d. The length s_1 increases for smaller inclination angles. If $\varphi \leq 30^\circ$, then $s_1 \geq 2.9 \text{ kpc}$, $q \leq 2$ and, since $s_1 < a$ and $R_{\text{CR}} \approx 1.2a$ according to numerical simulations and early type galaxy analyses, where a is the semi major axis of the bar, $R_{\text{CR}} \geq 3.5 \text{ kpc}$ (for $R_o = 8 \text{ kpc}$). Our simulations are presented in details in Fux (1998) and some mpeg movies, including live $\ell - V$ diagrams, are available at <http://obswww.unige.ch/~fux>.

References

- Dame, T.M, Ungerechts, H., Cohen, R.S. et al. 1987, ApJ 322, 706
 Fux, R. 1998, submitted to A&A




Figure 1. a) Observed CO l - V diagram (Dame et al. 1987), with the grey scale indicating the mean latitude of emission. b) Face-on spiral structure of an SPH model (bottom) and l - V traces of the associated arms (top). c) Parameters defining the position of the intersections of the lateral arms with the dustlane shocks. The location of the Sun is indicated by the $\odot$ symbol. d) Derived constraints on these parameters, with $q = s_1/s_2$ and where R_o is the galactocentric distance of the Sun.